

WARNINGS:

- Harmful if swallowed.
- Store away from food, feeds, seeds, fertilizers and other agricultural chemicals.
- Keep out of reach of children, animals and uninformed persons.

Although this product has been tested on a representative sample of economically important plant varieties under a large variety of soil and climatic conditions, the registration holder does not warrant that this product will be effective and safe to the crop under all conditions because action and effect of the chemical is affected by factors such as abnormal soil, climatic and storage conditions, as well as by the method and accuracy of application. The registration holder furthermore does not accept responsibility for either crop damage or non-performance due to failure to follow the label instructions or the occurrence of conditions which the registration holder could not reasonably have foreseen. Consult your supplier in the event of any uncertainty.

PRECAUTIONS:

- Do not inhale spray mist and/or fumes.
- In case of accidental contact with skin or eyes, wash immediately with plenty of water.
- Avoid spray drift onto other crops, grazing, rivers, dams, and any areas not under treatment.
- Thoroughly wash and rinse spray equipment after use and dispose of waste water where it will not contaminate crops, grazing, rivers and dams.
- Prevent contamination of food, foodstuffs, drinking water and eating utensils.
- Do not apply where the roots of susceptible plants can absorb the chemical.
- Rinse empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the contents of the spray tank before destroying the container by perforation and flattening and do not use any other purpose.

DIRECTIONS FOR USE:

- Use only as indicated.
- Shake well before use.

FARMAG DIURON 800 SC applied pre-emergence on its own is absorbed by the roots of germinating weeds and application must therefore be made before the weeds emerge on a cleanly cultivated soil. Rain or irrigation is required within one week after application to obtain good results. Application should preferably be made onto moist soil.

MIXING AND APPLICATION:

- Shake well before use.
- Mix the required quantity of **DIURON 800 SC** in a small quantity of water to make a smooth cream. Add this to the half filled spray tank and top up with water.
- Keep the chemical in continuous suspension in the spray tank by means of proper agitation.
- Apply in 200 – 400 litres water per hectare using flat fan type or flood jet nozzles.
- Accurately calibrate spray equipment prior to application.



FARMAG DIURON 800 SC

Reg No./Nr.: L 5663: Act/Wet 36 of/van 1947

A long acting residual suspension concentrate herbicide for the control of annual weeds as listed in citrus, avocados, bananas, coffee, mangoes, pineapples, macadamia and pecan nuts, and sugarcane, as well as in industrial areas along rail tracks and on road shoulders.

'n Suspensie konsentraat onkruidodder met lang nawerking vir die beheer van eenjarige onkruid soos aangetoon in sitrus, avokados, piesangs, koffie, mangoes, pynappels, makadamia en pekanneute en suikerriet, asook in nywerheidsgebiede langs spoorlyne en op padskouers.

HRAC HERBICIDE GROUP CODE	5	HRAC ONKRUIDDODER GROEP KODE
----------------------------------	----------	-------------------------------------

ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL:

Diuron (substituted urea).....800 g/lDiuron (gesubstitueerde ureum)

Manufacture Date:
Vervaardigingsdatum

Batch No.:
Lotnommer

NET CONTENTS/NETTO INHOUD

SEE ENCLOSED LEAFLET FOR FULL PARTICULARS
SIEN INGESLOTE VOUBILJET VIR VOLLE BESONDERHEDE

REGISTRATION HOLDER/REGISTRASIEHOUER:

CEDAR FALL PROPERTIES 130 (Pty) Ltd

Co. Reg. No./Mpy. Reg. Nr.: 2006/031383/07

61 Marshall Drive, Old Mill Industrial Park, Mount Edgecombe, South Africa, 4000 • Tel: 031 003 3486

UN No.: 3082



CAUTION
VERSIGTIG



DOSAGE RATE:

CROPS	ℓ/ha	COMMENTS
Citrus, avocados, mangoes, bananas (established orchards)	4,0	Apply onto clean cultivated soil during growing season. Repeat application after 2 – 3 months if required.
Pineapples (post plant or established plants)	4,0	Apply to clean cultivated moist soil. Split applications at half rates at 4 monthly intervals are preferable to a single application at full rate.
Coffee, macadamia and pecan nuts (established orchards only)	2,0	Apply to clean cultivated soil during Spring. Repeat applications after 2 – 3 months.
Plant sugarcane and ratoon crops	3,0 – 4,0	Apply as a uniform overall spray pre-plant or at the pre-emergence stage for weeds. Use higher rates for severe weed infestation or less susceptible weed.
Industrial use	10,0	Roadside, railways, industrial sites.

Some weeds species normally controlled by **DIURON 800 SC** include:

<i>Amaranthus hybridus</i>	Common pigweed
<i>Amaranthus spinosus</i>	Thorny pigweed
<i>Amaranthus thunbergii</i>	Red pigweed
<i>Bidens bipinnata</i>	Spanish blackjack
<i>Bidens pilosa</i>	Common blackjack
<i>Chenopodium album</i>	Fat hen
<i>Chenopodium carinatum</i>	Green goosefoot
<i>Cosmos bipinnatus</i>	Cosmos
<i>Digitaria sanguinalis</i>	Crab finger grass
<i>Eleusine indica</i>	Goose grass
<i>Gallinsoga parviflora</i>	Gallant Soldier
<i>Hibiscus trionum</i>	Kenaf
<i>Portulaca oleracea</i>	Pigweed
<i>Schkuhria pinnata</i>	Dwarf marigold
<i>Setaria verticillata</i>	Burr bristle grass
<i>Tagetes minuta</i>	Tall khaki weed



WAARSKUWINGS:

- Skadelik indien ingesluk.
- Bêre ver weg van voedsel, voer, sade, kunsmis en ander landboukundige chemikalieë.
- Hou buite bereik van kinders, oningeligte persone en diere.

Alhoewel hierdie produk getoets is op 'n verteenwoordigende monster van ekonomies belangrike plantsoorte onder 'n groot verskeidenheid van grond en klimaatstoestande, kan die registrasiehouer nie waarborg dat die produk effektief sal wees en veilig vir die gewas sal wees onder alle omstandighede, want aksie en gevolg van die chemiese produk word beïnvloed deur faktore soos abnormale grond, klimaat en bergingstoestande, sowel as die metode van toediening en die akkuraatheid daarvan. Die registrasiehouer aanvaar verdermeer ook nie verantwoordelikheid vir nog gewas beskadiging nog geen prestasie as gevolg van gebrek aan toepassing van die etiketinstruksies of die voorkoms van toestande wat nie voorsien kon word tydens registrasie nie. Raadpleeg die verskaffer in geval van enige onsekerheid.

VOORSORGMATREËLS:

- Moenie spuitnewel of walms inasem nie.
- In geval van toevallig kontak met vel of oë, was dit onmiddellik met baie water skoon.
- Vermy spuitdryf na ander gewasse, weivelde, riviere, damme en enige areas wat nie onder behandeling is nie.
- Was en spoel spuittoerusting deeglik skoon na gebruik, en raak ontslae van afvalwater waar dit nie ander gewasse, weivelde, riviere of damme sal besoedel nie.
- Verhoed besoedeling van voedsel, voer, drinkwater en eetgerei.
- Moenie toedien waar die wortels van die vatbare plante die chemikalieë kan absorber nie.
- Spoel die leë houer drie keer uit met 'n volume water wat gelykstaande is aan 'n minimum van 10% van die houer. Voeg die spoelwater by die inhoud van die spuittenk voordat die houer vernietig word deur gate daarin te maak en dit plat te trap en dit mag vir geen ander doeleindes gebruik word nie.

GERBUIKSAANWYSINGS

- Gebruik slegs soos voorgeskryf.
- Skud baie goed voor gebruik.

FARM AG DIURON 800 SC toegedien vooropkoms op sy eie, word geabsorbeer deur die wortels van die ontkiemende onkruid en daarom moet toediening gedoen word voordat die onkruid opkom op 'n skoon bewerkte stuk grond. Reën of besproeiing binne een week na toediening word verwag om die beste resultate te verkry. Toediening moet verkieslik gedoen word op vogtige grond.

MENG EN TOEDIENING

- Skud baie goed voor gebruik.
- Meng die voorgeskrewe hoeveelheid van **FARM AG DIURON 800 SC** in 'n klein hoeveelheid water om 'n gladde room te verkry. Voeg dit by die halfgevulde spuittenk en vul dit met water.


- Hou die chemikalieë onder voortdurende suspensie in die spuitenk deur middel van behoorlike beweging deur dit te roer.
- Dien toe in 200—400 liters water per hektaar deur gebruik te maak van 'n platwaaier tipe of 'n vloed spuitstraal spuitkoppe.
- Die spuittoerusting moet voor toediening korrek gekalibreer word.

DOSISSE

GEWASSE	ℓ/HA	OPMERKINGS
Sitrus, avokados, mangos, piesangs (gevestigde boorde).	4,0	Dien toe op skoon ontwikkelde grond gedurende die groeiseisoen. Herhaal toediening na 2 - 3 maande indien nodig.
Pynappels (na-planting of op gevestigde plante)	4,0	Dien toe op skoon ontwikkelde vogtige grond. Deel toediening in die halfdosis teen viermaandelikse intervalle is verkieslik bo 'n enkeltoediening teen die volle dosis.
Koffie, macadamia- en pekanneute (slegs gevestigde boorde)	2,0	Dien toe op skoon ontwikkelde grond gedurende Lente. Herhaal toediening na 2-3 maande.
Plant suikerriet en ratoen gewasse	3,0 - 4,0	Dien toe as 'n eenvormige algehele bespuiting tydens voorplant of op die vooropkoms stadium van onkruid. Gebruik hoër dosis vir erge onkruid infestasië of minder vatbare onkruid.
Industriële gebruik	10,0	Langs paaie, treinspore, industriële gebiede.

Sommige onkruidspesies wat normaalweg deur **FARM AG DIURON 800 SC** beheer word, sluit die volgende in:

<i>Amaranthus hybridus</i>	Gewone misbredie
<i>Amaranthus spinosus</i>	Doring misbredie
<i>Amaranthus thunbergii</i>	Rooi misbredie
<i>Bidens bipinnata</i>	Spaanse knapsekêrel
<i>Cosmos bipinnatus</i>	Kosmos
<i>Bidens pilosa</i>	Knapsekêrel
<i>Chenopodium album</i>	Wit hondebossie
<i>Chenopodium carinatum</i>	Groen hondebossie
<i>Digitaria sanguinalis</i>	Kruisvingergras
<i>Eleusine indica</i>	Osgras
<i>Gallinsoga parviflora</i>	Knokkruid
<i>Hibiscus trionum</i>	Terblansbossie
<i>Portulaca oleracea</i>	Porslein
<i>Schkuhria pinnata</i>	Klein kakiebos
<i>Setaria verticillata</i>	Klits borselgras
<i>Tagetes minuta</i>	Lang kakiebos





FARMAG DIURON 800 SC

Reg No./Nr.: L 5663: Act/Wet 36 of/van 1947

A long acting residual suspension concentrate herbicide for the control of annual weeds as listed in citrus, avocados, bananas, coffee, mangoes, pineapples, macadamia and pecan nuts, and sugarcane, as well as in industrial areas along rail tracks and on road shoulders.

'n Suspensie konsentraat onkruidodder met lang nawerking vir die beheer van eenjarige onkruid soos aangetoon in sitrus, avokados, piesangs, koffie, mangoes, pynappels, makadamia en pekanneute en suikerriet, asook in nywerheidsgebiede langs spoorlyne en op padskouers.

HRAC HERBICIDE GROUP CODE	5	HRAC ONKRUIDDODDER GROEP KODE
----------------------------------	----------	--------------------------------------

ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL:

Diuron (substituted urea).....800 g/lDiuron (gesubstitueerde ureum)

Manufacture Date:
Vervaardigingsdatum

Batch No.:
Lotnommer

NET CONTENTS/NETTO INHOUD

SEE ENCLOSED LEAFLET FOR FULL PARTICULARS
SIEN INGESLOTE VOUBILJET VIR VOLLE BESONDERHEDE

REGISTRATION HOLDER/REGISTRASIEHOUER:

CEDAR FALL PROPERTIES 130 (Pty) Ltd

Co. Reg. No./Mpy. Reg. Nr.: 2006/031383/07

61 Marshall Drive, Old Mill Industrial Park, Mount Edgecombe, South Africa, 4000 • Tel: 031 003 3486

UN No.: 3082



CAUTION
VERSIGTIG

